

A CONDENSED CONDENSATION PROOF OF A DETERMINANT EVALUATION CONJECTURED BY Greg KUPERBERG AND Jim PROPP

*Tewodros AMDEBERHAN*¹ and *Shalosh B. EKHAD*¹

Greg Kuperberg and Jim Propp [P] have conjectured the following determinant identity:

$$\det \left[\binom{i+j}{i} \binom{2n-i-j}{n-i} \right]_{0 \leq i, j \leq n} = \frac{(2n+1)^{n+1}}{(2n+1)!!} \quad ,$$

where $a!! := 0! \cdot 1! \cdot 2! \cdots a!$, and $a! := 1 \cdot 2 \cdots a$.

This is the special case ($m = n, a = b = 0$) of

$$\det \left[\binom{i+j+a+b}{i+a} \binom{2n-i-j-a-b}{n-i-a} \right]_{0 \leq i, j \leq m} = \frac{(a+b)!(2n+1)!^{m+1}(2n-m)!!m!!(m+a+b)!!(2n-m-a-b)!!a!!b!!(n-m-a-1)!!(n-m-b-1)!!}{a!b!(2n+1)!!(n-a)!!(n-b)!!(m+a)!!(m+b)!!(a+b)!!(2n-2m-a-b-1)!!} \quad ,$$

(*Rabbit*)

which follows immediately from Dodgson's[D] rule for evaluating determinants: (For any $n \times n$ matrix A , let $A_r(k, l)$ be the $r \times r$ connected submatrix whose upper leftmost corner is the entry $a_{k,l}$.)

$$\det A = \frac{\det A_{n-1}(1, 1) \det A_{n-1}(2, 2) - \det A_{n-1}(1, 2) \det A_{n-1}(2, 1)}{\det A_{n-2}(2, 2)} \quad . \quad (Lewis)$$

Indeed, let the left and right sides of (*Rabbit*) be $L_m(a, b)$ and $R_m(a, b)$ respectively. Dodgson's rule immediately implies that the recurrence:

$$X_m(a, b) = \frac{X_{m-1}(a, b)X_{m-1}(a+1, b+1) - X_{m-1}(a+1, b)X_{m-1}(a, b+1)}{X_{m-2}(a+1, b+1)} \quad ,$$

holds with $X = L$. Since $L_m(a, b) = R_m(a, b)$ for $m = 0, 1$ (check!), and the recurrence also holds with $X = R$ (check!²), it follows by induction that $L_m(a, b) = R_m(a, b)$ for *all* m . $\square$.

The present proof is in the spirit of [Z1]. Another proof can be found in [A1]. The same method yields a q-analog of (*Rabbit*), that can be found in [A2]. A beautiful combinatorial proof of (*Lewis*) can be found in [Z2]. An alternative proof of (*Rabbit*) is given in [K].

¹ Department of Mathematics, Temple University, Philadelphia, PA 19122, USA. E-mail: [tewodros,ekhad]@math.temple.edu; WWW: [http://www.math.temple.edu/~\[tewodros,ekhad\]](http://www.math.temple.edu/~[tewodros,ekhad]) .

² Divide both sides by the left, then use $r!/(r-1)!! = r!$ whenever possible, and then $r!/(r-1)!! = r$ whenever possible, reducing it to a completely routine polynomial identity. The small Maple package **rabbit** obtainable from our Home Pages, performs these steps mechanically.

References

- [A1] T. Amdeberhan, *A WZ proof of a determinant evaluation conjectured by Kuperberg and Propp*, exclusively published in Amdeberhan's Home Page <http://www.math.temple.edu/~tewodros>.
- [A2] T. Amdeberhan, *A q -generalization of a determinant evaluation conjectured by Kuperberg and Propp*, exclusively published in Amdeberhan's Home Page <http://www.math.temple.edu/~tewodros>.
- [D] C.L. Dodgson, *Condensation of Determinants*, Proceedings of the Royal Society of London **15**(1866), 150-155.
- [K] C. Krattenthaler, *E-mail message to T. Amdeberhan*, dated 9 Aug. 1996, 19 : 22 : 27 + 0100 (MET). By kind permission of Krattenthaler, it can be gleaned at the first author's Home Page <http://www.math.temple.edu/~tewodros>.
- [P] J. Propp, *E-mail message to D. Zeilberger*, dated 1 July 1996, 15:47:56 (EDT). By kind permission of Propp, it can be gleaned at the first author's Home Page <http://www.math.temple.edu/~tewodros>.
- [Z1] D. Zeilberger, *Reverend Charles to the aid of Major Percy and Fields Medalist Enrico*, Amer. Math. Monthly **103**(1996), 501-502.
- [Z2] D. Zeilberger, *Dodgson's determinant-evaluation rule proved by Two-Timing Men and Women*, to appear in the Elec. J. of Combinatorics [Wilf Festschrift volume]. It can be downloaded from Zeilberger's Home Page: <http://www.math.temple.edu/~zeilberg>.

Version of Sept. 4, 1996. First version: Aug. 2, 1996.